

Measures of Disease Frequency I

Seattle Epidemiology and
Biostatistics Summer Session
June, 2004

Prevalence and incidence

PREVALENCE is to INCIDENCE as...

Prevalence and incidence

PREVALENCE is to INCIDENCE as...

stock

is to

flow

Prevalence and incidence

PREVALENCE is to INCIDENCE as...

stock

is to

flow

wealth

is to

income

Prevalence and incidence

PREVALENCE is to INCIDENCE as...

stock is to flow

wealth is to income

account balance is to interest rate

Asymptomatic *C. trachomatis* infection

- Chronic state associated with ↑ risk of pelvic inflammatory disease, ectopic pregnancy, infertility in women

Asymptomatic *C. trachomatis* infection

- Chronic state associated with ↑ risk of pelvic inflammatory disease, ectopic pregnancy, infertility in women
- Curable with antibiotics

Asymptomatic *C. trachomatis* infection

- Chronic state associated with ↑ risk of pelvic inflammatory disease, ectopic pregnancy, infertility in women
- Curable with antibiotics
- 141,336 women tested upon entry into a national job training program during 1990–7

(Source: *Am J Public Health* 2001; 91:1287–90)

Asymptomatic *C. trachomatis* infection

- Chronic state associated with ↑ risk of pelvic inflammatory disease, ectopic pregnancy, infertility in women
- Curable with antibiotics
- 141,336 women tested upon entry into a national job training program during 1990–7
- 12.5% tested positive for *C. trachomatis*

(Source: *Am J Public Health* 2001; 91:1287–90)

Asymptomatic *C. trachomatis* infection

- Chronic state associated with ↑ risk of pelvic inflammatory disease, ectopic pregnancy, infertility in women
- Curable with antibiotics
- 141,336 women tested upon entry into a national job training program during 1990–7
- 12.5% tested positive for *C. trachomatis*
- What kind of disease frequency measure is this?

(Source: *Am J Public Health* 2001; 91:1287–90)

Prevalence

Prevalence

- Measures the frequency of being in the diseased *state* as of one point in time

Prevalence

- Measures the frequency of being in the diseased *state* as of one point in time
 - **Count of prevalent cases:** no. of people in the diseased state at that time

Prevalence

- Measures the frequency of being in the diseased *state* as of one point in time
 - **Count of prevalent cases:** no. of people in the diseased state at that time
 - **Prevalence:** $\frac{\text{No. of prevalent cases}}{\text{Size of population}}$


Prevalence

- Measures the frequency of being in the diseased *state* as of one point in time
 - **Count of prevalent cases:** no. of people in the diseased state at that time
 - **Prevalence:** $\frac{\text{No. of prevalent cases}}{\text{Size of population}}$
- A static measure: time is “frozen”


Prevalence

- Measures the frequency of being in the diseased *state* as of one point in time
 - **Count of prevalent cases:** no. of people in the diseased state at that time
 - **Prevalence:** $\frac{\text{No. of prevalent cases}}{\text{Size of population}}$
- A static measure: time is “frozen”
- Can be applied to any of several time scales:
 - Calendar time
 - Age
 - Time relative to some salient event

Prevalence diagram




Prevalence diagram




Point
in time

Prevalence diagram




$$\text{Prevalence} = 1/5 = 0.2 = 20\%$$

Chlamydia case detection over time



Newly recognized cases of infection were detected over the time period 1990–1997. So why isn't this *incidence* rather than *prevalence*?

Each person is *not monitored over time*



Common bile duct injury in cholecystectomy

- During gallbladder surgery, common bile duct may be injured

Common bile duct injury in cholecystectomy

- During gallbladder surgery, common bile duct may be injured
- Does frequency of injury depend on whether intraoperative cholangiography (IOC) used?

11

Common bile duct injury in cholecystectomy

- During gallbladder surgery, common bile duct may be injured
- Does frequency of injury depend on whether intraoperative cholangiography (IOC) used?

IOC used?

Yes	$2,380/613,706 = 0.39\%$
No	$5,531/956,655 = 0.58\%$

(Source: *JAMA* 2003; 289:1639–44)

11

Common bile duct injury in cholecystectomy

- During gallbladder surgery, common bile duct may be injured
- Does frequency of injury depend on whether intraoperative cholangiography (IOC) used?

IOC used?

Yes	$2,380/613,706 = 0.39\%$
No	$5,531/956,655 = 0.58\%$

- What kind of disease frequency measure is this?

(Source: *JAMA* 2003; 289:1639–44)

11

Common bile duct injury in cholecystectomy

- During gallbladder surgery, common bile duct may be injured
- Does frequency of injury depend on whether intraoperative cholangiography (IOC) used?

IOC used? **Cumulative incidence**

Yes	$2,380/613,706 = 0.39\%$
No	$5,531/956,655 = 0.58\%$

- What kind of disease frequency measure is this?

(Source: *JAMA* 2003; 289:1639–44)

11

Incidence

Incidence

- Measures frequency of disease *events* over time

Incidence

- Measures frequency of disease *events* over time
- Transition into diseased state may qualify as an event

Incidence

- Measures frequency of disease *events* over time
- Transition into diseased state may qualify as an event
- For some purposes, *count* of incident cases is enough

Incidence

- Measures frequency of disease *events* over time
- Transition into diseased state may qualify as an event
- For some purposes, *count* of incident cases is enough
- Otherwise, generally use either:
 - *Cumulative incidence*
 - *Incidence rate*

9

Cumulative incidence

- $$\frac{\text{No. of incident cases}}{\text{No. of people initially at risk}}$$

Cumulative incidence

- $$\frac{\text{No. of incident cases}}{\text{No. of people initially at risk}}$$
- Applies:
 - To a closed population initially at risk
 - When all persons are followed for same specified time period (at least approximately)


Cumulative incidence

- $$\frac{\text{No. of incident cases}}{\text{No. of people initially at risk}}$$
- Applies:
 - To a closed population initially at risk
 - When all persons are followed for same specified time period (at least approximately)
- A proportion or percentage


Cumulative incidence

- $\frac{\text{No. of incident cases}}{\text{No. of people initially at risk}}$
- Applies:
 - To a closed population initially at risk
 - When all persons are followed for same specified time period (at least approximately)
- A proportion or percentage
- Also called *attack rate* or *incidence proportion*

Cumulative incidence diagram



Cumulative incidence diagram



$$\text{Cumulative incidence} = 2/5 = 0.4 = 40\%$$

Incidence *rate*

- $$\frac{\text{No. of incident cases}}{\text{Amount of at-risk experience}}$$

Incidence *rate*

- $$\frac{\text{No. of incident cases}}{\text{Amount of at-risk experience}}$$
- Denominator usually expressed as person-time

Incidence *rate*

- $$\frac{\text{No. of incident cases}}{\text{Amount of at-risk experience}}$$
- Denominator usually expressed as person-time
- Can be applied:
 - To closed or open populations
 - When population members are at risk for different amounts of time

Incidence *rate*

- $$\frac{\text{No. of incident cases}}{\text{Amount of at-risk experience}}$$
- Denominator usually expressed as person-time
- Can be applied:
 - To closed or open populations
 - When population members are at risk for different amounts of time
- A rate, not a proportion: has units of 1/time

Incidence *rate*

- $$\frac{\text{No. of incident cases}}{\text{Amount of at-risk experience}}$$
- Denominator usually expressed as person-time
- Can be applied:
 - To closed or open populations
 - When population members are at risk for different amounts of time
- A rate, not a proportion: has units of 1/time
- Also called *incidence density* or *person-time incidence rate*

If detailed data available on each person

- Calculate each individual's contribution to numerator and denominator
 - Numerator: add up all qualifying disease events
 - Denominator: add up all time at risk


If detailed data available on each person

- Calculate each individual's contribution to numerator and denominator
 - Numerator: add up all qualifying disease events
 - Denominator: add up all time at risk
- Example: 15 needle-stick injuries in 1,500 nurse working hours = 1 case per 100 person-hours


If detailed data available on each person

- Calculate each individual's contribution to numerator and denominator
 - Numerator: add up all qualifying disease events
 - Denominator: add up all time at risk
- Example: 15 needle-stick injuries in 1,500 nurse working hours = 1 case per 100 person-hours
- Calculations may differ depending on whether recurrent events in same person qualify


If recurrent events *do* qualify...




If recurrent events *do* qualify...




If recurrent events *do* qualify...




If only *first* events qualify...



If only *first* events qualify...



If only *first* events qualify...



If detailed data *not* available on each person

$$\frac{\text{No. of incident cases}}{\left(\text{Avg. size of population at risk} \right) \times \left(\text{Duration of observation period} \right)}$$

If detailed data *not* available on each person

$$\frac{\text{No. of incident cases}}{\left(\begin{array}{c} \text{Avg. size of} \\ \text{population at risk} \end{array} \right) \times \left(\begin{array}{c} \text{Duration of} \\ \text{observation period} \end{array} \right)}$$

Example: 100 heart attack cases occur in 1 year
in a city of 100,000 persons

$$\begin{aligned} \text{Incidence rate} &\approx \frac{100 \text{ cases}}{100,000 \text{ persons} \times 1 \text{ year}} \\ &\approx 1.0 \text{ cases per 1000 person-years} \end{aligned}$$

Variants of incidence

- Mortality
 - Measures burden of fatal disease *on the population as a whole*

Variants of incidence

- Mortality
 - Measures burden of fatal disease *on the population as a whole*
 - Cumulative mortality = $\frac{\text{No. of fatal cases}}{\text{No. initially at risk}}$

Variants of incidence

- Mortality
 - Measures burden of fatal disease *on the population as a whole*
 - Cumulative mortality = $\frac{\text{No. of fatal cases}}{\text{No. initially at risk}}$
 - Mortality rate = $\frac{\text{No. of fatal cases}}{\text{Person-time at risk}}$

Variants of incidence

- Mortality
 - Measures burden of fatal disease *on the population as a whole*
 - Cumulative mortality = $\frac{\text{No. of fatal cases}}{\text{No. initially at risk}}$
 - Mortality rate = $\frac{\text{No. of fatal cases}}{\text{Person-time at risk}}$
- Case fatality
 - Measures risk that a person *who gets the disease* will die of it

Variants of incidence

- Mortality
 - Measures burden of fatal disease *on the population as a whole*
 - Cumulative mortality = $\frac{\text{No. of fatal cases}}{\text{No. initially at risk}}$
 - Mortality rate = $\frac{\text{No. of fatal cases}}{\text{Person-time at risk}}$
- Case fatality
 - Measures risk that a person *who gets the disease* will die of it
 - Case fatality = $\frac{\text{No. of fatal cases}}{\text{Total no. of cases}}$

Proportional mortality

- $$\frac{\text{Deaths from a certain disease}}{\text{Total no. of deaths}}$$

Proportional mortality

- $$\frac{\text{Deaths from a certain disease}}{\text{Total no. of deaths}}$$
- Denominator only a proxy for true population at risk


Proportional mortality

- $$\frac{\text{Deaths from a certain disease}}{\text{Total no. of deaths}}$$
- Denominator only a proxy for true population at risk
- Sometimes used when:
 - Comparing deaths from 2+ diseases in same population


Proportional mortality

- $$\frac{\text{Deaths from a certain disease}}{\text{Total no. of deaths}}$$
- Denominator only a proxy for true population at risk
- Sometimes used when:
 - Comparing deaths from 2+ diseases in same population
 - Comparing frequency of death from a certain disease across 2+ populations of unknown size

Pitfall of proportional mortality—1



Pitfall of proportional mortality—1



Deaths:

Disease X

20


30

Other causes

80

170

Pitfall of proportional mortality—1



Pitfall of proportional mortality—2

Village	Deaths in 1 year		Proportional mortality
	Total	Disease X	
A	100	20	20%
B	200	30	15%

Pitfall of proportional mortality—2

Village	Deaths in 1 year		Proportional mortality	Pop. size
	Total	Disease X		
A	100	20	20%	1,000
B	200	30	15%	1,000

Pitfall of proportional mortality—2

Village	Deaths in 1 year		Proportional mortality	Pop. size	Mortality rate*
	Total	Disease X			
A	100	20	20%	1,000	20
B	200	30	15%	1,000	30

*Deaths per 1,000 person-years

Pitfall of proportional mortality—2

Village	Deaths in 1 year		Proportional mortality	Pop. size	Mortality rate*
	Total	Disease X			
A	100	20	20%	1,000	20
B	200	30	15%	1,000	30

*Deaths per 1,000 person-years

- Proportional mortality greater in Village A

Pitfall of proportional mortality—2

Village	Deaths in 1 year		Proportional mortality	Pop. size	Mortality rate*
	Total	Disease X			
A	100	20	20%	1,000	20
B	200	30	15%	1,000	30

*Deaths per 1,000 person-years


- Proportional mortality greater in Village A
- True mortality rate greater in Village B (!)

Period prevalence

- $$\frac{\text{No. of pre-existing cases} + \text{No. of new cases}}{\text{Total population}}$$


Period prevalence

- $$\frac{\text{No. of pre-existing cases} + \text{No. of new cases}}{\text{Total population}}$$



Period prevalence


- $$\frac{\text{No. of pre-existing cases} + \text{No. of new cases}}{\text{Total population}}$$



Period prevalence = $4/5 = 0.8$

Period prevalence

- $$\frac{\text{No. of pre-existing cases} + \text{No. of new cases}}{\text{Total population}}$$




Period prevalence = $4/5 = 0.8$

- Really a hybrid of prevalence and cumulative incidence

Period prevalence

- $$\frac{\text{No. of pre-existing cases} + \text{No. of new cases}}{\text{Total population}}$$




Period prevalence = $4/5 = 0.8$

- Really a hybrid of prevalence and cumulative incidence
- Interpretable only for closed population

Period prevalence

- $$\frac{\text{No. of pre-existing cases} + \text{No. of new cases}}{\text{Total population}}$$



Period prevalence = $4/5 = 0.8$

- Really a hybrid of prevalence and cumulative incidence
- Interpretable only for closed population
- Example: period prevalence of diabetes in pregnant mothers is about 25.3 per 1,000

Introduction to Epidemiologic Methods — Summer, 2004
Discussion Questions: Measures of Disease Frequency—1

1. Between January, 1990 and March, 2002, the Food and Drug Administration received 3,339 reports of rhabdomyolysis, a rare disease that involves massive breakdown of skeletal muscle cells, among users of a class of drugs called statins, which reduce blood cholesterol. Some 1,899 of the rhabdomyolysis cases had taken a particular drug, cerivastatin. During that time period, cerivastatin accounted for about 2.0% of all statin prescriptions.

Assume for present purposes that the number of days' worth of medication dispensed with each prescription was about the same for all statins and that the reporting of rhabdomyolysis was complete. From the data given, can you estimate the ratio of the incidence of rhabdomyolysis among cerivastatin users to the incidence of rhabdomyolysis among users of other statins? If so, what is it? If not, why not?

2. Hepatitis C is of concern to the Department of Veterans Affairs (VA) because many VA users are thought to fall into groups at high risk for this disease. It is believed to cause cirrhosis, liver cancer, and other complications. Among the early studies that have sought to gauge the extent of infection among VA users were:
 - In early 1994, 839 consecutive patients admitted to the Washington, D.C. VA Medical center were tested, and 173 (20.6%) of them were found to be infected with hepatitis C.
 - On March 17, 1999, some 26,000 outpatients at VA clinics nationwide who agreed to be tested were screened for hepatitis C. The exact number of positives is not available, but the prevalence was reported to be 8–10%

What main concern(s) would you have about using these results as estimates of the prevalence of infection among veterans? Would you expect the estimates to be biased upward or downward?